

# ACARSTIN FLO®

Registration No. L4165 Act 36/1947

Residual contact suspension concentrate acaricide for the control of Two-spotted Spider Mite (Red Spider) (including organophosphate resistant mites) and European Red Mite on apple, pear, plum, peach and nectarine trees, tomatoes and hops.

Active Ingredient:  
Cyhexatin (Organotin)

600 g / ℥

Net Volume

1 ℥

Aktiewe Bestanddeel:  
Siheksatin (Organotin)

Netto Volume

Registered by:  
Sipcam Southern Africa (Pty) Ltd  
Co. Reg. No. 1996/009324/07  
P.O. Box 516, Suider Paarl 7624  
Tel. No. (021) 874 1055



Distributed by:  
Chempack (Pty) Ltd  
Co. Reg. No. 2002/007885/07  
P.O. Box 516, Suider Paarl 7624  
Tel. No. (021) 874 1055

**chempac**  
progressive agri-care

Acarstin Flo® is the registered trade mark of  
SIPCAM S.p.A., Milan, Italy

Acarstin Flo® is die geregistreerde handelsmerk van  
SIPCAM S.p.A., Milan, Italië

Batch number

Lotnommer

Date formulated

Datum geformuleer

UN NUMBER 3020



HARMFUL  
SKADELIK



VN NOMMER 3020

## WARNINGS

DO NOT APPLY WITHIN 3 DAYS OF HARVEST OF APPLES, PEARS, TOMATOES AND HOPS; 7 DAYS ON PLUMS, PEACHES AND NECTARINES.

Poisonous if swallowed. May cause irritation of the eyes and mucous membranes. Toxic to fish and wild life. Keep out of reach of children, un informed persons and animals. Store away from food and feeds.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS, QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE PEST AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

## PRECAUTIONS

Do not inhale spray mist. When handling the concentrate, wear gloves and a face shield. Wash with soap and water after use. Avoid contact with eyes, skin and clothing. In case of skin or eye contact, wash skin thoroughly with soap and water and immediately flush eyes with running water. Do not eat drink or smoke whilst mixing and applying or before washing hands and face. Remove contaminated clothing and wash before re-use. Avoid drift of spray onto other crops, grazing, rivers or dams. Clean applicator after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams. Prevent contamination of food, feedstuffs, drinking water and eating utensils. Rinse empty container three times with a volume of water equal to at least one tenth of that of the container. Add the rinsings to the contents of the spray tank before destroying the empty container. Do not use empty container for any other purpose.

## DIRECTIONS FOR USE

### Compatibility:

Acarstin Flo is compatible with most commonly used pesticides but should not be used together with alkaline materials, lime sulphur or sulphur.

### Mixing and spraying instructions:

Mix the required amount of Acarstin Flo with a little water before adding the final volume of water. Spray thoroughly to wet all parts of the tree or plant. Commence spraying as soon as an increase in the number of mites is noticed. Repeat as necessary.

CROP/PEST	DOSAGE	REMARKS
Apples, Pears, Plums, Peaches, Nectarines Two-spotted spider Mite and ERM (European Red Mite)	50 mL / 100L water (1,5 – 2,5 L / ha)	Apply as a high volume spray. Dosage per hectare depends on the size of trees.
Tomatoes Two-spotted Spider Mite	625 – 750 mL / ha	Apply as a full cover spray when infestation is noticed. Repeat when necessary. A second application is recommended in case of severe infestation to ensure good control. The dosage depends on plant size.
Hops Two-spotted Spider Mite	50 mL / 100L water	Apply as a full cover spray when infestation is noticed. Repeat when necessary. A second application is recommended in case of severe infestation to ensure good control.