

STRIKER

Reg. No. L6628 Act. No. 36 of 1947

A water dispersible granular formulation of sulphur for use as a fungicide and acaricide to control pests on the following crops: avocado, citrus, vines, deciduous fruits, papaya, vegetables and ornamentals.

Reg. Nr. L6628 Wet Nr. 36 van 1947

'n Water dispergeerbare korrelformulasie van swael vir gebruik as 'n swam- en mytdoder vir die beheer van peste op die volgende gewasse: avokado, sitrus, druwe, sagtevrugte, papaja, groente en sierplante.

Active ingredient:
Sulphur

800 g/kg

Aktiewe bestanddeel:
Swael

Net Contents

25 kg

Netto inhoud

Manufactured by :
QUIMETAL INDUSTRIAL S.A.
Santa Ester 748, San Miguel, Santiago, CHILE

Vervaardig deur :
QUIMETAL INDUSTRIAL S.A.
Santa Ester 748, San Miguel, Santiago, CHILE

Registered and Distributed by:
CHEMPAC (Pty) Ltd. Reg. No 2002/007885/07
P O Box 516, Suider Paarl 7624
Tel.No. (021) 874 1055

Geregistreer en Versprei deur:
CHEMPAC (Edms) Bpk. Reg. Nr. 2002/007885/07
Posbus 516, Suider Paarl 7624
Tel.Nr (021) 874 1055

Batch No.

Lot Nr.

Date of Manufacture

Datum van Vervaardiging



CAUTION
VERSIGTIG



WARNINGS:

- Do not apply to grapevines later than four to six weeks before harvest.
- May irritate the eyes, skin and respiratory tract.
- Avoid spraying during the hottest part of the day in intense sunshine.
- Flammable – do not store near open flame or sparks.
- Keep out of reach of children, uninformed persons and animals.
- Store in cool, dry places away from food and feed.

PRECAUTIONS:

- Do not inhale dust or spray mist. Wear rubber gloves and face shield when mixing.
- Wash with soap and water after use.
- Do not eat, drink or smoke while mixing or spraying, or before washing hands and face.
- Prevent contamination of food, eating utensils and drinking water.
- Prevent spray drift onto other crops, grazing, rivers, dams and areas not under treatment.
- Empty bags should be disposed of by burying or burning. When burning ensure bags are totally empty, bearing in mind the flammable nature of the product and the possible consequences of burning product. Do not use empty bags for any other purpose.
- Clean application equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

DIRECTIONS FOR USE: Use only as directed.

MIXING AND SPRAYING INSTRUCTIONS:

Half fill the spray tank with water. Mix the required amount of STRIKER separately with a small amount of water. Add this to the water in the spray tank and stir well. Fill spray tank with water to the required volume. Maintain agitation throughout spraying operation.

CROP	PEST	DOSAGE/ 100 ℥ WATER	REMARKS
Apples and Pears	Bud Mite	400 g (10,0 – 14,0 kg/ha)	Full cover spray applied at closed bud stage.
	Scab and Powdery Mildew	HV. 200 – 400 g (3,0 – 14,0 kg/ha)	<i>Pre-blossom application:</i> 300 – 400 g/100ℓ water. <i>Early post-blossom application:</i> 250 g/100ℓ water. <i>Late post-blossom application:</i> 200 g /100 ℥ water Do not use on sulphur-shy cultivars.
Avocados	Thrips	300 g	Apply when pest is noticed
Beans	Red Spider Mite	400 g	Apply as a full cover spray every 10 – 14 days.
Citrus	Flat Mite	150 g	Apply during December to May when pest is noticed.
	Grey Mite	200 g	Apply after petal drop. Repeat two months later.
	Rust Mite	300g	Apply when first signs of shadow like blemishes are noticed on fruit during December – May.
	Silver Mite	250 g	Apply in spring if russetting occurred in previous season.
	Thrips	300 g	Apply first spray at 100% petal drop. Repeat 3 times at 10-day intervals if necessary. Allow 4 weeks between oil and sulphur treatments.
Cucurbits	Powdery Mildew	300 g	Apply as soon as disease is noticed. Repeat every 1 – 2 weeks as long as weather favours disease. Use up to 1500 ℥ spray mixture / ha depending on foliage. Do not use on muskmelon and cucumbers.
Grapes	Erinose Mite	HV. 150 – 300 g	Dosage depends on severity of disease or pest. Heavy sprays of sulphur may cause burn especially on hot days.
	Powdery Mildew	HV. 150 – 300 g	Apply when shoots are 10 cm long with a second application at flowering and a third in about the third week of December. <i>Winter rainfall region:</i> Start application as soon as shoots are 10 cm long and repeat every three weeks for as long as necessary. Will also control bud mite (Erinose mite) <i>Summer rainfall region:</i> Start application in mid summer especially if it is very dry or when first symptoms appear.
Ornamental Flowers	Red Spider Mite	400 g	Apply as soon as pest is noticed. Repeat if necessary. Full cover spray.
	Powdery Mildew	300 g	Full cover sprays every 10 – 14 days depending on weather favourable for disease. Roses: Apply weekly as long as new growth is formed.
	Thrips	400 g	Apply weekly when pest damage is noticed. Sulphur may injure some ornamentals.
Papaya	Powdery Mildew	300 g	Apply as soon as disease is noticed. Apply weekly when disease is prevalent.
Peaches	Powdery Mildew	HV. 150 – 300 g	Apply as soon as first symptoms appear and repeat every 10 – 14 days as long as weather favours the disease. Dosage depends on severity of disease.
Peas	Powdery Mildew	300 g	Apply when first symptoms of disease appear and repeat every 7 – 10 days. Apply up to 1 000 ℥ spray mixture /ha depending on plant size.
Plums, Apricots and Peaches	Rust	HV. 150 – 300 g	Dosage depends on severity of disease. <i>Winter rainfall region:</i> Start applications early November. <i>Summer rainfall region:</i> Apply from mid-February onwards as soon as symptoms are noticed. Repeat every 10 – 14 days.
Tomatoes	Rust Mite Erinose Mite	400 g	Apply as soon as the pest is noticed. Repeat at 2 – 4 week intervals as necessary.

=====

WAARSKUWING:

- Moenie later as vier tot ses weke voor oes van tafel- of wyndriuw gebruik nie.
- Mag oë, vel en asemhalingstelsel irriteer.
- Moenie gedurende die warmste deel van die dag in direkte sonlig spuit nie.
- Vlambaar: Moet nie naby oop vlamme of vonke geberg word nie.
- Hou buite bereik van kinders, oningesigte persone en diere.
- Bêre in 'n koel, droë plek weg van voedsel en veevoer.

VOORSORGMAATREEËLS:

- Moenie stof of spuitmis inasem nie. Dra rubberhandskoene en 'n gesigskerm wanneer gemeng word.
- Was met seep en water na gebruik.
- Moet nie eet, drink of rook wanneer gemeng of gespuit word, alvorens hande en gesig gewas is nie.
- Voorkom besoedeling van voedsel, eetgerei en drinkwater.
- Voorkom die wegdrywing van spuitnewel na ander gewasse, weiding, riviere, damme en ander oppervlakte wat nie behandel word nie.
- Leë verpakking moet begrawe of verbrand word. Wanneer dit verbrand word, maak seker dat dit heeltemal leeg is. Neem in ag dat die produk vlambaar is en wees bedag op die moontlike gevolge van verbranding van die produk. Moet nie leë sakke vir enige ander doel gebruik nie.
- Maak spuitapparaat skoon na gebruik en raak ontslae van waswater waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

MENG EN TOEDIENINGS INSTRUKSIES:

Vul spuittenk half vol met water. Meng die nodige hoeveelheid STRIKER apart met 'n klein hoeveelheid water. Voeg by die water in spuittenk en roer goed. Vul die spuittenk met water tot die verlangde volume. Spuitmengsel in spuitenk moet voortdurend geroer word tydens bespuiting.

GEWAS	PLAAG	DOSIS/100 ℥ WATER	OPMERKINGS
Appels en Pere	Knopmyt Skurf en Poeieragtige skimmel	400 g (10,0 – 14,0 kg/ha) HV. 200 – 400 g (3,0 – 14,0 kg/ha)	Volle dektoediening met toeknopstadium. <i>Voor bloei-toediening:</i> 300 – 400 g/100 ℥ water. <i>Vroeë nabloei-toediening:</i> 250 g/100 ℥ water. <i>Laat na-bloei-toediening:</i> 200 g/100 ℥ water. Moet nie op swaelsku kultivars gebruik nie.
Avokado's	Blaaspootjie	300 g	Spuit sodra plaag opgemerk word.
Boontjies	Rooispinmyt	400 g	Dien toe as 'n volle bedekking elke 10 – 14 dae.
Druwe	Erinosemyt Poeieragtige Skimmel	HV. 150 – 300 g HV. 150 – 300 g	Dosis hang af van erns van siekte of plaag. Swaar toedienings van swael kan wingerde brand veral op warm dae. Begin met toediening sodra die lote 10 cm lank is opgevolg met 'n tweede toediening met blomtyd en 'n derde toediening in die derde week van Desember. <i>Winterreënvalstreek :</i> Begin met toediening sodra lote 10 cm lank is en herhaal elke 3 weke solank dit nodig is. Sal ook knopmyt beheer. <i>Somerreënvalstreek :</i> Dien toe vanaf middel somer of wanneer eerste tekens van die siekte opgemerk word, veral as dit baie droog is.
Ertjies	Poeieragtige Skimmel	300 g	Dien toe wanneer die eerste tekens van die siekte opgemerk word en herhaal elke 7 – 10 dae. Dien tot 1000ℓ spuitmengsel/ha toe, afhangende van plantgrootte.
Pampoengewasse	Poeieragtige Skimmel	300 g	Dien toe sodra die siekte opgemerk word. Herhaal elke 1 – 2 weke vir solank die weerstoestande die siekte begunstig. Gebruik tot 1500 ℥ spuitmengsel/ha afhangende van plantgrootte. Moenie op spanspek en komkommers gebruik nie.
Papaja	Poeieragtige Skimmel	300 g	Dien toe sodra eerste simptome verskyn. Dien weekliks toe solank siekte voorkom.
Perskes	Poeieragtige Skimmel	HV.150 – 300 g	Dien toe sodra eerste simptome verskyn. Herhaal elke 10 – 14 dae solank weerstoestande die siekte begunstig. Dosis hang af van die erns van die siekte.

Pruime, Appelkose en Perskes	Bruinroes	HV. 150 – 300 g	Dosis hang af van erns van die siekte. <i>Winterreënvalstreek:</i> Begin toediening vroeg in November. <i>Somerreënvalstreek:</i> Dien toe vanaf middel Februarie of wanneer eerste tekens van die siekte opgemerk word. Herhaal elke 10 – 14 dae.
Sierplante - Blomme	Rooispinmyt	400 g	Dien toe sodra plaag opgemerk word. Herhaal indien nodig. Volle dektoediening.
	Poeieragtige Skimmel	300 g	Volle dektoediening elke 10 – 14 dae solank as wat die weer gunstig vir die siekte bly. Rose: Spuit elke week solank as nuwe groei gevorm word.
	Blaaspootjie	400 g	Dien weekliks toe wanneer plaagskade opgemerk word. Swael mag sekere plante beskadig.
Sitrus	Platmyt Grysmyt Roesmyt	150 g 200 g 300 g	Dien toe in Desember tot Mei sodra plaag opgemerk word. Dien toe na blomblaarval. Herhaal twee maande later. Dien toe sodra die eerste tekens van skaduagtige verkleuring op die vrugte opgemerk word in Desember tot Mei.
	Silwermyt	250 g	Dien toe gedurende lente indien skilverruwing die vorige seisoen voorgekom het.
	Blaaspootjie	300 g	Eerste toediening met 100% blomblaarval. Herhaal 3 keer met 10 dae tussenposes indien nodig. Laat 4 weke tussen olie en swael behandelings toe.
Tamaties	Roesmyt Erinosemyt	400 g	Dien toe sodra plaag opgemerk word. Herhaal met 2 – 4 weke tussenposes indien nodig.